













# ID Phyt Capcin recommendations



| Species / physiological stage   |   | Minimum           | Practical  | Maximum  | Duration / Application                                       | Matrix values |
|---|---|-------------------|--|--|--|---------------|
|  <p><b>Ruminant</b></p>  |  <p>Dairy cows</p>                           | 600 mg/cow/day    | 800 mg/cow/day   | 1000 mg/cow/day                                      | Min. 45 days<br>If possible, start a few days before calving | Yes           |
|   |  <p>Beef cattle</p>                          | 100 mg/animal/day | 300 mg/animal/day  | 500 mg/animal/day<br>(acidogenic diets, heat stress) | Min. 30 days   | Yes           |
|  <p><b>Swine</b></p>     |  <p>Lactating sows</p>                       |                   | 200 g/t  |  | If possible, start during late-gestation                     |               |
|   |  <p>Piglets &amp; growing pig up to 25kg</p> |                   | 200 g/t  |  |  |               |
|   |  <p>Growing pig 25-70 kg</p>                 |                   | 150 g/t  |  |  |               |
|   |  <p>Growing pig 70-120 kg</p>               |                   | 100 g/t  |  |  |               |
|  <p><b>Poultry</b></p> |  <p>Broilers</p>                           | 100 g/t           | 200 g/t (starter)<br>150 g/t (grower)<br>100g/t (finisher)   | 200 g/t  |  | Yes           |
|   |  <p>Layers</p>                             | 100 g/t           | 100 g/t  | 125-150 g/t<br>(Heat stress)                         |  |               |
|   |  <p>Turkey</p>                             | 100 g/t           | 200 g/t up to 8-10 weeks<br>150 g/t up to 12 weeks<br>100 g/t up to 14-16 weeks<br>0 or 100 g/t > 16 weeks |  |  |               |